

Diversity of *Striga gesnerioides* on cowpea in Ghana

Mr Frederick Awuku

MSc Plant Breeding and Genetics, University of Ghana.

Supervisors: Professor Pangirayi Tongoona and Dr Agyemang Danqah, University of Ghana, and Dr Francis Kusi, CSIR-SARI.



Mr Frederick Awuku

Project overview

Striga gesnerioides represents a major constraint over cowpea yield in northern Ghana but the identity and diversity of races has not been established. This project sought to characterize the distribution and diversity of *S. gesnerioides* in Ghana. The parasite was found to occur over a large area of the northern part of the country. Both purple and white flowered types are present, although the former predominate by about 3:1. The reaction of 27 cowpea varieties to eight *S. gesnerioides* populations sampled from three separate regions of northern Ghana revealed that some of the material was resistant to all of the populations, some was susceptible to all of the populations, and some showed a differential response. The existence of the differential response implied the presence of more than one race of *S. gesnerioides* in Ghana. A genotypic survey of 27 samples of *S. gesnerioides* based on 73 newly developed PCR markers revealed two main clusters, supporting the suggestion that multiple strains of the parasite are present in Ghana. The identities of the races present in Ghana are likely SG3, SG2 and SG5.



Map of Ghana indicating the *Striga gesnerioides* collection sites for this project (i); flower colour variation in *S. gesnerioides* sampled (ii-v).